

## Effect of fermentation periods of poultry manure on its characteristics when fermented with different ways

Nabaa O. Alburih<sup>id</sup> and Zainab K. Hasan<sup>id</sup>

Soil Science and Water resources, College Agriculture, Basrah University, Iraq

\*Corresponding author's e-mail: [nabaaottman@gmail.com](mailto:nabaaottman@gmail.com); [zinabkadm006@gmail.com](mailto:zinabkadm006@gmail.com)

Following the scientific management of organic manure (poultry) may contribute to reducing environmental pollution and human health, as the current study was conducted with the aim of recycling poultry manure accumulated in the field of poultry farming at the research station - Faculty of Agriculture - Karma site and the impact of varying fermentation durations on organic manure, specifically animal (poultry manure) fermented by different fermentation methods and its impact on the rest of the factors to produce highly decomposed organic manure free of pathogens. Several traditional fermentation methods (compost and water immersion) were followed. and modern (bioreactor), and for a fermentation period of 90 days with an estimate of most of the qualitative characteristics of the organic manure produced and within periods of 15 - 30 - 45 - 60 - 75 - 90 days. The results of the experiment of incubation of poultry manure showed that all the studied traits have changed with the progress of fermentation time and in the same direction, the results demonstrated that fermentation in the bioreactor unit has given the highest values of total nitrogen, total phosphorus and potassium 32.53, 16.72, 20.86, g.kg<sup>-1</sup> for the elements mentioned sequentially, while the lowest values of pH 6.68 and C/N ratio 9.96 compared to other fermentation methods.

**Keywords:** Bioreactor, Compost, Soaking in Water, Poultry Manure, Manure. Organic manure management, Pathogen-free organic manure, Fermentation period, Environmental pollution reduction.

### INTRODUCTION

The treatment of soil with organic residues from different sources (animal - plant - and others) is an agricultural practice that began in ancient times and with the beginning of human knowledge of agriculture (Al-Fares, 2017), as organic and animal matter, in particular, has a role in improving the chemical, physical and biological characteristics of the soil in addition to fertility, in addition to that organic matter as added manures may be fully or partially complementary to the traditional mineral recommendation of nitrogen, phosphate and other manures, which are often exposed to several obstacles and problems and in most

Soils that lead to the loss of the largest part of them, which limits or reduces the maximum productivity of crops (Ren *et al.*, 2020; Cai *et al.*, 2021; Tang *et al.*, 2021). The process of recycling (Recycling) animal manure (such as poultry, for example) is one of the promising scientific methods that support soil fertility and crop productivity, on two sides, The first is to find a safe way to get rid of the accumulations of these residues and their negative impact on the environment and human health, and the second hand is to improve the

quality and specifications of the effective unit of the nutritional value of those residues (Ali *et al.*, 2014; Duncan, 2005; Escobar and Hue, 2008). Especially when treated as organic manure free of pathogens, bush seeds, and others (Saveyn and Eder, (2014)). The use of organic manure (compost) decomposed air method is beneficial in improving the content of soil of organic matter as this method is one of the means of biological treatment (bio) of organic manure, whether of plant or animal origin by beneficial bacteria that work to decompose these materials when appropriate environment of moisture (60) % and optimum temperature (60 – 70) We get a good organic manure that works to enrich the soil with living organisms when added to it and install Atmospheric air nitrogen and dissolving phosphorus and potassium, so it is easy for the plant that absorbs it regularly, so it grows with high efficiency (Al-Madini, 2012). Water immersion is a normal practice used by field farmers in Basra Governorate that benefits the soil (Abdulkareem, 2010). With the rapid development of modern agriculture, the use of anaerobic organic manure fermentation was resorted to in the reactor unit. Biological, which is a source of clean renewable energy (renewable energy) and the use of biotechnology

(biogas) or what is known as a bioreactor is one of its main outputs is an organic manure with high-quality specifications that differs in its characteristics from organic manure prepared by traditional methods, including compost and immersion in water (Arnos, 2022; Zubaidi, 2020; Jurgutis *et al.*, 2020; Zhang, 2021; Amjid *et al.*, 2011). The results of recent research studies confirmed by Arnous (2022), Zubaidi (2020), Jurgutis *et al.* (2020), and Amjid *et al.* (2011). The fermentation process is affected by several factors including temperature, starter addition, continuous stirring, pH, C/N, C/P, and fermentation duration. This study aimed to recycle (Recycling) poultry manure accumulated in the field of breeding Poultry and the extent of the effect of the period on the fermentation of organic manure, specifically animal (poultry manure) fermented by different fermentation methods (anaerobic fermentation in the bioreactor unit, compost fermentation, and water immersion fermentation) and its impact on the rest of the factors.

## MATERIALS AND METHODS

**Sample Collection:** Poultry manure was collected from the poultry breeding field of the College of Agriculture - the University of Basra, the site of Karma Ali, mixed well, and dried in the oven at a temperature of 50 °C, then ground and sieved from a sieve with a capacity of 1 mm openings and estimated for its primary characteristics as shown in Table 1.

**Table 1. Some characteristics of raw poultry manure before fermentation.**

Characteristics	Value	Unite
pH (5:1)	7.20	-
EC (5:1)	19.6	dS m <sup>-1</sup>
Organic matter	769.20	g kg <sup>-1</sup>
Organic carbon	447.20	
Total -N	35.70	
Total-P	17.15	
Total-K	19.27	
C/N	12.53	-
C/P	26.07	-
PW	80.53	(w:w)

### Experimental Setup:

1. Fermented soft poultry manure (Table 1) by anaerobic fermentation in the locally manufactured bioreactor and established and designed from a previous study at the research station of the College of Agriculture - University of Basra - Karma Ali, foreign substances were removed and mixed with water and by 250 kg Manure: 500 liters of water in addition to its moisture content of 80% Taking into account the mixing to make the mixture homogeneous with providing it with a starter and after the stability of the basic manure characteristics of anaerobically fermented manure such as nitrogen and the

ratio C: N Especially after 90 days of fermentation, samples were taken for post-fermentation analysis as in Table 2.

2. Poultry manure taken from the poultry field was immersed in plastic tanks filled with water prepared in a way that allows gas exchange between the perforated pipes embedded in the tanks and the surrounding atmosphere, taking into account the maintenance of a constant water level inside the tank throughout the fermentation period.
3. The composting method is summarized in preparing the ground by digging a hole with dimensions (length \* width 1 \* 1 m, depth 1.5 m), including three layers The first layer was placed 8-10 cm of fronds and placed on top of it a layer of soft poultry manure and a thickness of 2 cm and covered with a layer of soil to a depth of 3 cm and with the succession of the mentioned layers with mixing for homogeneity, then add a certain amount of urea manure. Moistened with R.O. water by 60%. The pile was covered with plastic wrap, taking into account the need to stir every three days for ventilation during the fermentation process, which took 3 months.

**Table 2. Some Characteristics of fermented poultry manure for 90 days in different ways**

Characteristics	Unite	Compost	Soaking in water	Biogas
pH(5:1)	-	7.02	7.20	6.92
Ec(5:1)	dS m <sup>-1</sup>	19.86	24.10	10.97
organic matter	g kg <sup>-1</sup>	445.90	567.90	557.30
organic carbon		259.24	330.17	324.01
total -N		25.90	29.57	32.53
total-P		14.89	16.07	16.72
total-K		18.90	19.37	20.86
C/N	-	10.09	11.16	9.96
C/P	-	17.41	20.54	19.37

### Analytical Methods

**pH:** Potting in a suspension of organic manure: water (5:1) using a pH meter. Liquid manure (separated from the bioreactor) has been estimated directly using a pH meter, as reported by Page *et al.* (1982).

**EC:** Estimated in an organic manure suspension: water (5:1) using Ec-meter while liquid manure (separated from the bioreactor) was estimated directly using Ec-meter, as reported in Page *et al.* (1982).

**Total nitrogen:** Estimated in steam distillation digestion solution as reported in Bremner (1970).

**Total phosphorus:** It was estimated in the digestion solution after modifying the acidity of the mixture, and using a color spectrometer at a wavelength of 700 nm in the blue color method as reported in Murphy and Riley (1962).

**Total potassium:** Estimated in the digestion solution using a flame photometer after adjusting the acidity of the solutions



according to [Page et al. \(1982\)](#).


**Organic matter:** It was estimated according to the loss-on-ignition (LOI) method, where the model was burned at 550° C for 8 hours as reported in [ECS \(1999\)](#) from which organic carbon was calculated.

**C/N Ratio nitrogen:** It was calculated from the quotient of organic carbon divided by total

**C/P Ratio:** Calculated from the quotient of organic carbon divided by total phosphorus.

## RESULTS AND DISCUSSION

**pH :** The results of Fig. 1 showed that there is a significant effect ( $P<0.05$ ) of fermentation periods for fermented poultry manure in different ways, specifically by immersion in water and bioreactor methods in the pH of fermented manure, as the pH of fermented manure decreased by water immersion and bioreactor methods from 7.2 (beginning of fermentation) to 6.5 and 4.6 within 15 days for two methods mentioned sequentially, while the pH of the manure prepared by the traditional compost method to 8, as it was noted that regardless of the method in which the poultry manure was fermented, the microbial activity is the main factor causing the change in the characteristics of the subject materials, including their pH ([Abu Hmaid, 2017](#)). It is worth noting that in both methods of water immersion and bioreactor may have optional anaerobic microorganisms (water immersion) and compulsory anaerobic (bioreactor) in The possibility of converting the fermentation medium to acidic, especially the beginning of fermentation, because the developing anaerobic bacteria work to break down easily degradable materials and convert them into Organic acids in the pH phase ([Macias-Corral et al. 2008](#)).



F: Fermentation method T: Time


**Figure 1. Effect of fermentation duration on the pH of fermented organic manure by different methods**

The results of Fig. 1 showed that there was a slight rise in the pH of fermented manure in the bioreactor unit over 75 days, as the pH of the fermented manure in this way did not exceed

6 compared to the other two methods (immersion in water and compost), which indicates a state of stability and a specialized type of compulsory anaerobic microorganisms according to the biological system, which is consistent with ([Mao et al., 2015](#)).

It was shown that the pH index of anaerobically fermented materials decreases from 7 (beginning of fermentation) to 6.8-5.2 (end of anaerobic fermentation), attributing this to the predominance of what is known as hydrolysis and Acidogenesis, especially when combined with methane-releasing bacteria. Methanogens drop pH from 7.2 to 5.3 due to the breakdown of organic fatty acids (VFA) ([Getachew, 2012](#)).

**Total organic carbon:** The results of Fig. 2 showed that there is a significant effect ( $P<0.05$ ) in reducing the organic carbon of fermented poultry manure by different methods by increasing the fermentation time compared to the period of 0 days and for all fermentation methods as the results showed that all fermentation methods followed the same behavior in decreasing their carbon content with the progress of fermentation time, the decrease was clear in the treatment of anaerobic fermentation in the bioreactor unit compared to compost and immersion in water and all fermentation periods may be because the carbon in the manures Anaerobic fermentation in the bioreactor unit is more easily biodegradable than the carbon in aerobically fermented fertilize RS ([Odlare et al. 2011](#)). This also came in agreement with [Al-Jumaili and Al-Azzawi \(2016\)](#) who obtained the continued decrease in the percentage of organic matter in anaerobic conditions compared to aerobic fermentation and pointed out that the difference in decline between anaerobic fermentation and aerobic fermentation increases in breadth with the passage of fermentation periods, attributing the reason for this to the availability of appropriate environmental conditions for the growth The proliferation of microorganisms is mainly responsible for degradation and therefore a continuous decrease in organic carbon.




F: Fermentation method T: Time



**Figure 2. Effect of fermentation duration on the total organic carbon content of fermented organic manure by different methods.**


The results of Fig. 3 showed the effect of fermentation time on the nitrogen content of poultry manure prepared in different ways, as it is clear from the results that the percentage of nitrogen decreased during the first 15 days and for all fermentation methods compared to the period of 0 days due to the consumption of nitrogen microorganisms and then took to increase their nitrogen content and have followed the same behavior in all fermentation methods, and the method affected significantly the nitrogen content and it was the highest value of nitrogen content in poultry manure prepared anaerobically in the bioreactor unit compared to other methods For all fermentation periods, the highest nitrogen value was 32.53 g kg<sup>-1</sup> at 90 days of fermentation in the bioreactor unit compared to compost and water immersion, at which the values reached 25.90 and 29.57 g kg<sup>-1</sup> for the methods mentioned sequentially, these results were identical to what was obtained [Abdulkareem\(2010\)](#) for the products of the decomposition of poultry manure prepared in three ways (air, anaerobic and water immersion). This may be due to the decrease in the compost pile after fermentation and the disintegration of the raw materials involved in the composition of these residues as a result of the loss of easily degradable compounds and the release of CO<sub>2</sub>, thus increasing the release of nitrogen at the expense of losing weight in the compost pile ([Orrico et al. 2012](#)). It is expected that nitrogen loss in the form of ammonia gas will occur in all fermentation methods used in the study except for the anaerobic fermentation method, these results were similar to what happened to [Jacob et al. \(1995\)](#), who proved that the rapid collection of manure and fermentation under anaerobic conditions far from the encouraging effects of loss cases will lead to maintaining the high nutritional value of organic manure.



F: Fermentation method T: Time

**Figure 3. Effect of fermentation duration on the total nitrogen content of fermented organic manure by different methods.**

**Phosphorus content:** The phosphorus content in the manure prepared in different ways increased with the increase in the period of fermentation, specifically after the first 15 days of fermentation, where all methods followed the same behavior (Figure 4) and reached the highest phosphorus content after 80 days of fermentation This result is consistent with many studies, which indicated that the element phosphorus is an essential element in the process of producing organic manure and proved an increase in the percentage of phosphorus in the final yield of organic manure and attributed this increase to the decomposition of organic residues During the fermentation process and the release of this element from fermented residues due to the activity of microorganisms in addition to the decrease in the unit weight in the compost pile with time ([Samaha, 2018](#)). The loss of dry matter and the decrease in unit weight as a result of the fermentation of organic residues, regardless of the method, leads to an increase in its nutrient content ([Yagodin, 1984](#)). The decomposition products (Figure 8) under anaerobic conditions (bioreactor) showed the highest significant amount of phosphorus, which amounted to 16.72 g kg<sup>-1</sup> compared to compost and immersion in water 14.88 and 16.06 g kg<sup>-1</sup> These results were consistent with [Al-Assafi et al. \(2010\)](#), and also consistent with [Abed et al. \(2015\)](#) who found that the highest movement of phosphorus liberation reaches after 80-120 days and that the concentration of phosphorus in anaerobically fermented residues is higher than aerobic fermented residues with an increase of 57.67% and 45.99% of the content of Total phosphorus in the mixture and for the methods mentioned sequentially.




F: Fermentation method T: Time

**Figure 4. Effect of fermentation duration on the total phosphorus content of fermented organic manure by different methods.**



**Potassium content:** The results of Fig. 5 showed the effect of fermentation time on the potassium content of poultry residues prepared in different ways. It is noted from the figure that the potassium content in fermented residues in different ways increased significantly with time and this increase was associated with what the conditions and period of fermentation did in reducing the weight of dry matter for soft animal manures (Ashour, 1995), which led to an increase in potassium content in fermented poultry. The results showed that the treatment of anaerobic fermented manure (bioreactor) was significantly superior to the rest of the fermentation methods (compost and immersion with water) and all fermentation periods, as the highest value of potassium content in the products of the fermentation process reached 20.86 g kg<sup>-1</sup> for anaerobic fermented manure, while it reached 18.90 and 19.37 g kg<sup>-1</sup> for the mentioned methods sequentially compared to the period of 0 days, which amounted to 19.27 g kg<sup>-1</sup>. It is also seen from the figure that the overlap between time and method was not significant in the potassium content of poultry manure.




F: Fermentation method T: Time

**Figure 5. Effect of fermentation duration on the total potassium content of fermented organic manure by different methods.**

**C/N ratio:** The C/N ratio is used as an indicator of the rate of degradation, where a decrease in the C/N ratio or a narrowing of the value between nitrogen and carbon indicates that there has been significant decomposition (Al-Assafi *et al.*, 2010). Fig. 6 shows the difference in the C/N ratio. For organic residues prepared in different ways over time, the results obtained showed that increasing the fermentation period leads to a decrease in the percentage of C/N significantly and all fermentation methods have followed the same behavior by

giving them a gradual decrease significantly with the progress of the fermentation period. The reason for the decrease in the percentage of C/N over time is due to the decrease in the percentage of carbon during the decomposition and increase in the percentage of total nitrogen. These results are consistent with the findings of Abdel-Hadi (2008) and Al-Fadhli (2011) and Šubová *et al.*, (2021). The results also showed significant differences between the fermentation methods at the end of decomposition and the superiority of anaerobic fermented manure by giving the lowest C/N ratio, which amounted to 9.96 compared to compost and water immersion, which amounted to 10.02 and 11.16 respectively, they proved Yan *et al.* (2021) that the C/N ratio of 8.84 is the optimal ratio for manure produced from bioreactor units, and this ratio is considered to have a stronger effect in promoting corn growth. soil moisturizing and availability of biology. Awasthi *et al.* (2014) showed that a C/N ratio equal to or less than 25 is an indicator of manure maturity, while Iqbal *et al.* (2015) indicated that C/N less than 20 is evidence of acceptable organic manure and if it drops to 15 or less, it is more acceptable.



F: Fermentation method T: Time


**Figure 6. Effect of fermentation duration on C/N ratio of fermented organic manure by different methods**

**C/P ratio:** Fig. 7 shows the effect of fermentation time on the ratio of C/P in poultry manure prepared in different ways, as the effect of time significantly on the values of c/p for poultry manure and notes a gradual decrease significantly in the ratio of C/P in all fermentation methods during the decomposition process with time as a result of the high rate of decomposition of C and the increase in phosphorus concentration (Islam, 2021), as the release of phosphorus increases with the increase in the duration of fermentation (Figure 8), and the method also significantly affected the values of C/P and the





values reached 17.44, 20.55 and 19.37 for each of the compost, water immersion, and bioreactor sequentially.



F: Fermentation method T: Time:

**Figure 7. Effect of fermentation duration on C/P ratio of fermented organic manure by different methods.**

**Conclusions:** 1- The results of the incubation experiment showed that all the studied traits have changed with the duration of fermentation and in the same direction, and the results showed that fermentation in the bioreactor unit has given the highest values of total nitrogen, total phosphorus, and potassium 32.53, 16.72, 20.86, g kg<sup>-1</sup> for the elements mentioned sequentially, while the lowest values of pH 6.68 and C/N ratio 9.96

2- The process of anaerobic fermentation with a unit and the bioreactor and the establishment of this group does not negatively affect the environment at all stages of construction and production and supports sustainable agriculture through environmental preservation and recycling of organic manure and can be established in most circumstances because it does not need complications in raw materials and implementation 3- Fermentation of poultry manure prepared in different ways for 90 days gave the best results for the basic elements

**Recommendations:** 1- Exploiting animal manure, especially abandoned poultry manure, by converting it into high-quality organic manure free of pathogens, reducing the percentage of unpleasant odors, and working to clean the environment to contribute to sustainable agriculture and achieve a return for agricultural crop Continuity of research in this field and the work of a developed study to maintain all conditions of the device

2- The need to provide a bioreactor device for each farm supported by a separation device to establish an integrated unit and project with an economic return for the agricultural crop

**Authors' contributions:** Authors' contributions: Nabaa O. Alburih<sup>1</sup>, Zainab K. Hasan: conceived and designed the experiments, performed the experiments, analyzed the data. and wrote the paper, reviewed the manuscript. All authors read and approved the final manuscript.

**Funding:** by Authors

**Ethical statement:** This article does not contain any studies with human participants or animal performed by any of the authors.

**Availability of data and material:** We declare that the submitted manuscript is our work, which has not been published before and is not currently being considered for publication elsewhere

**Code Availability:** Not applicable

**Consent to participate:** All authors are participating in this research study.

**Consent for publication:** All authors are giving the consent to publish this research article in JGIAS

**Acknowledgements:** This work is a part of the Research in College of Agriculture, University of Basrah, Iraq.

## REFERENCES

- Abdel-Hadi, M. A., and A. El-Azeem. 2008. Effect of heating, mixing, and digester type on biogas production from buffalo dung. *Misr Journal of Agricultural Engineering* 25:1454-1477.
- Abdulkareem, M. A. 2010. Effect of composting methods of manures on arylsulfatase activity in sandy soil. *Al-Mustanstriya J Sci* 21:1-10.
- Abed, A. A., B.Hassan and H. E. Nouri. 2015. The effect of fermentation method and processing with phosphate rock in recycling solid organic manure for homes into organic manures. *Anbar University Journal of Pure Sciences*. M pp.
- Abu Hmaid, D. M. 2017. Wheat straw treatment with base and microwave radiation for the production of biogas by anaerobic fermentation method (Doctoral dissertation, Islamic University of Gaza).
- Al-Assafi, A. A., Jamal, S., and H. Bardan, 2010. Recycling and treatment of agricultural solid manure for cities. *Iraqi Journal of Desert Studies* 5:1-6
- Al-Fadhli, J. T. M. 2011. The effect of organic and mineral fertilizing on the growth and yield of potatoes. (. *(Solanum tuberosum L)*). Ph.D. thesis, University of Baghdad.
- Al-Fares, M. A. N. 2017. Design, manufacture, and evaluation of the performance of soil tillage machines at different depths and the addition of organic manure and its impact on some soil characteristics and flower plant yield. Doctoral thesis. The University of Basra. Faculty of Agriculture.
- Ali, N. S., S. R. Hamdallah and A. A. Shaker. 2014. Soil fertility. Dar Al-Kutub Al-Alamia for Printing, Publishing, and Distribution. Amman, Jordan. GSD.307.
- Al-Jumaili, A. A. and S. S. Al-Azzawi. 2016. Organic carbon and active carbon are associated with some soil characteristics. *Al-Qadisiyah Journal of Agricultural Sciences* 61-7.



- Al-Madini, A. B. M. 2012. The effect of adding compost on the chemical and fertility qualities of the soil, College of Agricultural and Food Sciences / King Faisal University
- Amjid, S. S. M. Q. Bilal, M. S. Nazir and A. Hussain. 2011. Biogas is a renewable energy resource for Pakistan. *Renewable and Sustainable Energy Reviews* 15:2833-2837.
- Arnous, A. M. J. 2022. Barley plant response to liquid organic manure produced from anaerobic fermentation of poultry manure. Master Thesis, College of Agriculture - University of Basrah - Iraq.
- Ashour, N. I. 1995. The effect of storage period and methods on the chemical composition of soft organic manure Basrah Journal of Agricultural Sciences 8:1-8.
- Awasthi, M. K., A. K. Pandey, J. Khan, P. S. Bundela, J. W. C. Wong and A. Selvam. 2014. Evaluation of thermophilic fungal consortium for organic municipal solid manure composting. *Bioresour. Technol* 168:214-221.
- Bremner, J. M. 1970 . Regular Kjeldahl methods. In: A.L. Page; R.H. Miller and D. R. Keeney .1982 (eds.) *Methods of soil analysis. Part 2, 2<sup>nd</sup> Ed.* ASA. Inc. Madison, Wisconsin, U.S.A.
- Cai, F., P. Luo, J. Yang, M. Irfan, S. Zhang, N. An. 2021. Effect of long-term fertilization on ammonia-oxidizing microorganisms and nitrification in brown soil of northeast China. *Front. Microbiol.* 11. doi: 10.3389/fmicb.2020.622454.
- Duncan, I. J. H. 2005. Science-based assessment of animal welfare: Farm animals. *Review Sci. Techni. Office Inter. des Epi.* 24:483-492.
- ECS. European Committee for Standardisation. 1999. soil improvers and growing media - determination of organic matter content and ash technical committee CEN/TC 223.
- Escobar, M.E.O. and N.V. Hue. 2008. Temporal changes of selected chemical characteristics in three manure-Amended soils of Hawaii. *Bioresour* 99:8649-8654.
- Getachew, D. 2012. Effect of adding urea on biogas production potentials of selected fruit manures in Addis Ababa, Ethiopia.
- Iqbal, M. K., A. Nadeem, F. Sherazi, and R. A. Khan. 2015. Optimization of process parameters for kitchen manure composting by response surface methodology. *International Journal of Science, Environment and Technology.* 12:1759-1768.
- Islam, M. R., S. Bilkis, T. S. Hoque, S. Uddin, M. Jahiruddin, M. M. Rahman and R. Datta. 2021. Mineralization of Farm Manures and Slurries under Aerobic and Anaerobic Conditions for Subsequent Release of Phosphorus and Sulphur in Soil. *Sustainability* 13: 8605.
- Jurgutis, L., A. Slepeliene, J. Volungevicius and K. Amaleviciute-Volunge. 2020. Biogas production from chicken manure at different organic loading rates in a mesophilic full-scale anaerobic digestion plant. *Biomass and Bioenergy* 141: 105693
- Macias-Corral, M.; Samani, Z.; Hanson, A.; Smith, G.; Funk, P.; Yu, H. and J. Longworth. 2008. Anaerobic digestion of municipal solid manure and agricultural manure and the effect of co-digestion with dairy cow manure. *Bioresource technology* 99: 8288-8293.
- Mao, C., Y. Feng, X. Wang and G. Ren. 2015. Review on research achievements of biogas from anaerobic digestion. *Renewable and sustainable energy reviews.* 45: 540-555.
- Murphy, T. and J. R. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta.* 27:31-36.
- Odlare, M., V. Arthurson, M. Pell, K. Svensson, E. Nehrenheim and J. Abubaker 2011. Land application of organic manure — effects on the soil ecosystem. *Appl. Energy* 88:2210-2218.  
<http://dx.doi.org/10.1016/j.apenergy.2010.12.043>.
- Orrico, A. C. A., Centurion, S. R., Farias, R. M. D., Orrico Junior, M. A. P., & R. G. Garcia. 2012. Effect of different substrates on composting of poultry litter. *Revista Brasileira de Zootecnia* 41:1764-1768.
- Page, A. L.; Miller, R. H. and D. R. Keeney, 1982. *Methods of soil analysis. Part (2) 2nd Agronomy* 9.
- Ren, T., X. Yu, J. Liao, Du, Y. Zhu, L. Jin. 2020. The application of biogas slurry rather than biochar increases soil microbial functional gene signal intensity and diversity in a poplar plantation. *Soil Biology and Biochemistry.* 146, 107825: doi: 10.1016/j.soilbio.2020.107825.
- Samaha, A. Y. 2018. Compost resulting from pruning residues and its impact on the productivity of some varieties of peaches and apples in the Zabadani area. Master's Thesis. Damascus University. Faculty of Agriculture, Syrian Arab Republic.
- Saveyn, H. and P. Eder. 2014. End-of-manure criteria for biodegradable manure subjected to biological treatment (compost and digestate): tech. propo. Report EUR, 264251. Sevilla, Spain.
- Šubová, E., N. Sasáková, F. Zigo, I. Mindžáková, M. Vargová, J. Kachnič and K. V. Laktičová, 2021. Amendment of livestock manure with natural zeolite-clinoptilolite and its effect on decomposition processes during composting. *Agriculture* 11 980.
- Tang, Y., Luo, L., Carswell, A., Misselbrook, T., Shen, J., and J. Han, 2021. Changes in soil organic carbon status and microbial community structure following biogas slurry application in a wheat-rice rotation. *Science of the Total Environment* 757:143786. doi: 10.1016/j.scitotenv.2020.143786.
- Yagodin, B. A. 1984. *Agricultural chemistry. Part 2.* Mir pub. Moscow.



- Yan, L., C. Liu, Y. Zhang, S. Liu, and Y. Zhang. 2021. Effects of C/N ratio variation in swine biogas slurry on soil dissolved organic matter: Content and fluorescence characteristics. *Ecotoxicology and Environmental Safety* 209:111804.
- Yaqoub, I. B., I. A. Nawal and H. J. Tamimi. 1995. The effect of storage period and methods on the chemical characteristics of organic manure. *Basra Journal of Agricultural Sciences* 8:81-97.
- Zhang, H., S. Li, X. Zheng, J. Zhang, N. Bai, H. Zhang and W. Lv. 2021. Effects of biogas slurry combined with chemical manure on soil bacterial and fungal community composition in a paddy field. *Frontiers in Microbiology* 2338
- Zubaidi, B. M. J. 2020. Effect of Interaction between Organic Fertilization and Biogas-Treated Water Quality and Output on Some Soil Characteristics and Potato Yield. Doctoral thesis. Faculty of Agriculture. Muthana University. Iraq.

